
## 6.13 Improper Integrals

Write your questions and thoughts here!

Improper integrals are integrals with infinite limits of integration or have an infinite discontinuity on the interval.

If f(x) is continuous on  $[a, \infty)$ , then  $\int_a^{\infty} f(x) dx =$ 

If f(x) is continuous on  $(-\infty, b]$ , then  $\int_{-\infty}^{b} f(x) dx =$ Provided the limits exist!



If the limit exists, the improper integral is said to \_\_\_\_\_ . If the limit does not exist, the integral is said to \_\_\_\_\_. 2.  $\int_{1}^{\infty} \frac{1}{x} dx$ 



Improper *p*-integral:  $\int_{1}^{\infty} \frac{1}{x^{p}} dx$  converges if

and diverges if

Write your questions and thoughts here! Remember the definite integral  $\int_{a}^{b} f(x) dx$ , requires the interval to be finite and the FTC requires that f(x) be continuous on [a, b]. If the integral does not meet these requirements, we may need to manipulate the problem.

Another form of the Improper Integral is  $\int_{-\infty}^{\infty} f(x) dx$ , with f(x) continuous on  $(-\infty, \infty)$ . Let x = c be any real number in the interval  $(-\infty, \infty)$ , then

$$\int_{-\infty}^{\infty} f(x) \, dx = \int_{-\infty}^{c} f(x) \, dx + \int_{c}^{\infty} f(x) \, dx.$$

(It's easiest to use 0 here for c). If either of these integrals diverge, then the whole diverges.

3. 
$$\int_{-\infty}^{\infty} e^x \, dx$$

If f(x) is continuous on [a, b) and has an infinite discontinuity at b, then

$$\int_{a}^{b} f(x) \, dx =$$

If f(x) is continuous on (a, b] and has an infinite discontinuity at a, then  $\int_{a}^{b} f(x) dx =$ 

$$4. \quad \int_0^2 \frac{x+2}{\sqrt{x^2+4x}} dx$$

If f(x) is continuous on the interval [a, b], except for some c in (a, b) at which f has an infinite discontinuity, then  $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$ . 5.  $\int_{-1}^1 \frac{1}{x} dx$ 

## 6.13 Improper Integrals

Calculus Evaluate each integral.		Practice
1. $\int_{1}^{\infty} \frac{1}{x^2} dx$	2. $\int_0^\infty \frac{2}{x^2 + 4x + 3} dx$	
$3. \int_0^1 \frac{x+1}{\sqrt{x^2+2x}} dx$	$4. \int_{1}^{\infty} x e^{-x} dx$	

$5. \int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$	6. $\int_{-1}^{0} \frac{1}{x^5} dx$
7. $\int_0^\infty e^{-x} dx$	<sup>9</sup> Determine all the values of <i>n</i> for which $\int_{-1}^{1} dx$
7. J <sub>0</sub> e ux	8. Determine all the values of p for which $\int_0^1 \frac{1}{x^p} dx$ converges.

## 6.13 Improper Integrals

9. If g is a twice-differentiable function, where g(2) = 1 and  $\lim_{x \to \infty} g(x) = 8$ , then  $\int_2^{\infty} g'(x) dx$  is

10. If *R* is the unbounded region between the graph of  $y = \frac{x}{(1+x^2)^2}$  and the *x*-axis for  $x \ge 0$ , then the area of *R* is

## A) -1 (B) 0 (C) $\frac{1}{2}$ (D) infinite

11. For what values of p will  $\int_1^\infty \frac{1}{x^{7p-3}} dx$  converge?

A) 
$$p < 0$$
 (B)  $p > 0$  (C)  $p > \frac{4}{7}$  (D)  $p < \frac{4}{7}$